arXiv:1408.3729v2 [math.GT] 7 Apr 2015

THE COMPLETE LIST OF PRIME KNOTS WHOSE FLAT PLUMBING BASKET NUMBERS ARE 6 OR LESS

YOON-HO CHOI, YUN KI CHUNG, AND DONGSEOK KIM

Abstract. Flat plumbing basket surfaces of links were introduced to study the geometry of the complement of the links. These flat plumbing basket surface can be presented by a sequential presentation known as flat plumbing basket code first found by Furihata, Hirasawa and Kobayashi. The minimum number of flat plumbings to obtain a flat plumbing basket surfaces of a link is defined to be *the flat plumbing basket number* of the given link. In present article, we use these sequential presentations to find the complete classification theorem of prime knots whose flat plumbing basket number 6 or less. As applications, this result improves the work of Hirose and Nakashima which finds the flat plumbing basket number of prime knots up to 9 crossings.

Seifert surfaces, banded surfaces, flat plumbing basket surfaces, flat plumbing basket number

[2000]57M25, 57M27

1. Introduction

Orientable surfaces whose boundary is the given link, known as *Seifert surfaces* have been studied for many interesting invariants of links such as Seifert pairings, Alexander polynomials, signatures and etc. A *plumbing surface* obtained from a 2-dimensional disc by plumbings annuli found by Rudolph [17] used to study extensively for the fibreness of links and surfaces [2–4, 7, 14, 16, 19]. In particular, if we only use flat annuli plumbings, the resulting surface is called a *flat plumbing surface*. The main focus of the present article is flat plumbing basket surfaces, a precise definition can be found in Definition 2.1. A flat plumbing basket surface can be regarded as a flat plumbing surface, but not vice versa. There exists a Seifert surface which is obtained from a disk by successively plumbing flat annuli, but which is not isotopic to any flat plumbing basket surface [2].

The third author's first preprint about these plumbing surfaces from a canonical Seifert surface had a critical mistake. In the process of resolving this mistake, the third author, Kwon and Lee proved the existence of banded surfaces and flat banded surfaces [13] by weakening some conditions of plumbings. The third author also proved that every link L is the boundary of an oriented surface which is obtained from a graph embedding of a dipole graph, this surface is also known as a braidzel surface [15], and a complete bipartite graph $K_{2,n}$, where all voltage assignments on the edges of dipole graph and $K_{2,n}$ are 0 [11]. The mistake was finally fixed in [10].

2000 *Mathematics Subject Classification.* 57M25, 57M27.

Key words and phrases. Flat plumbing basket surfaces, Seifert surfaces, banded surfaces.

The present work is one of articles in this series of results presenting links as a boundary of the surface obtained in a embedding of certain graphs as described in [6]. One might consider these plumbing surfaces as special embeddings of the bouquets of circles [5].

A sequence of articles by Furihata, Hirasawa and Kobayashi [2] and the third author [10] proved the existence of a flat plumbing basket surface of a given link L. We can define the *flat plumbing basket number* of L, denoted by $fpbk(L)$, to be the minimal number of flat annuli to obtain a flat plumbing basket surface of the link L. However, finding the flat plumbing basket number of a link L is very difficult and has been beyond the reach for more than 15 years from the first invention by Rudolph [17]. because it is defined to be the minimum over all possible flat plumbing basket surfaces whose boundaries is the given link L.

The work of Furihata *et al.* [2] provided not only the existence theorem using a very tangible alternating definition of the flat plumbing basket surface but also a coding algorithm, the resulting code is called *flat plumbing basket code*, to present links as the boundaries of flat plumbing basket surfaces from a special closed braid presentation of the link.

In present article, we use these sequential codes to find all prime knots of the flat plumbing basket number 6 in Theorem 3.6 by applying DT-code and a computer program "knotfinder" and "knotscape" of `Knotscape` [20].

When the third author first presented this work at the TAPU conference in 2013 summer, Carter pointed out that this flat plumbing basket code of a link can be very useful to calculate Alexander polynomials because all components in Seifert matrix can be found directly from the presentation and they are either 0 or ± 1. A very recent work by Hirose and Nakashima [8] found a theorem which provide two lower bounds of the flat plumbing basket number using Alexander polynomials and genera of links. Using these lower bounds, they succeed to find the flat plumbing basket number of all prime knots up to 9 crossings except 24 knots.

As an application of our classification theorem, we find the flat plumbing basket number of five knots out of 24 knots and sharpens the range of the flat plumbing basket number of three knots.

The outline of this paper is as follows. We first provide some preliminary definitions and results in Section 2. We provided an explicit coding algorithm to find the flat plumbing basket presentation of a link from its braid presentation and canonical Seifert surface. Also we provide two classification theorems of the flat plumbing basket number of 4 and 6 with a explanation how we find DT-code and use the computer program "knotfinder" of `Knotscape` in Section 3. We conclude with a remark on further research in Section 4.

2. Preliminaries

A compact orientable surface $\mathcal{F}$ is called a *Seifert surface* of a link L if the boundary of $\mathcal{F}$ is isotopic to the given link L. The existence of such a surface was first proven by Seifert using an algorithm on a diagram of L, this algorithm was named after him



FIGURE 1. (a) A geometric shape of α, B_α and C_α on a Seifert surface S and (b) a new Seifert surface $\overline{S}$ obtained from S by a top A_0 plumbing along the path α.

as *Seifert's algorithm* [18]. A Seifert surface $\mathcal{F}_L$ of an oriented link L produced by applying Seifert's algorithm to a link diagram is called a *canonical Seifert surface*.

The main topic of the article is the flat plumbing basket surfaces. Rudolph [17] first defined the top plumbing as follows. Let α be a proper arc on a Seifert surface S. Let B_α be a 3-cell on the positive side of the oriented surface S along a tubular neighborhood C_α of α on S. Let $A_n \subset B_\alpha$ be an n times full twisted annulus such that $A_n \cap \partial B_\alpha = C_\alpha$. The *top plumbing* on S along a path α is the new surface $S' = S \cup C_\alpha$ where A_n, B_α, C_α satisfy the previous conditions as depicted in Fig. 1. Thus, two consecutive plumbings are non-commutative in general. Rudolph found a few interesting results with regards to the top and bottom plumbings in [17]. For the rest of article, all plumbings are top plumbing unless state differently.

Definition 2.1. *A Seifert surface $\mathcal{F}$ is a* flat plumbing basket surface *if $\mathcal{F} = D_2$ or if $\mathcal{F} = \mathcal{F}_0 *_\alpha A_0$ which can be constructed by plumbing A_0 to a flat plumbing basket surface $\mathcal{F}_0$ along a proper arc $\alpha \subset D_2 \subset \mathcal{F}_0$. We say that a link L admits a* flat plumbing basket representation *if there exists a flat plumbing basket surface $\mathcal{F}$ such that $\partial\mathcal{F}$ is equivalent to L.*

An alternative definition of the flat plumbing basket surfaces is given in [2] and it is very easy to follow. The *trivial open book decomposition* of $\mathbb{R}^3$ is a decomposition of $\mathbb{R}^3$ into the half planes in the following form. In a cylindrical coordinate, it can be presented

$$\mathbb{R}^3 = \bigcup_{\theta\in[0,2\pi)} \{(r,\theta,z)|r \geq 0, z \in \mathbb{R}\}$$

where $\{(r,\theta,z)|r \geq 0, z \in \mathbb{R}\}$ is called a *page* for $\theta \in [0, 2\pi)$. Let $\mathcal{O}$ be the *trivial open book decomposition* of the 3-sphere $\mathbb{S}^3$ which is obtained from the trivial open book decomposition of $\mathbb{R}^3$ by the one point compactification. A Seifert surface is said to be a flat plumbing basket surface if it consists of a single page of $\mathcal{O}$ as a 2-disc D^2 and finitely many bands which are embedded in distinct pages [2]. Flat plumbing basket surfaces of (i) the trefoil knot and (ii) the figure eight knot in the trivial open book decomposition are depicted in Fig. 2 where D^2 is presented as a



FIGURE 2. (a) A flat 4-banded surface of the trefoil knot and (b) a flat 4-banded surface of the figure eight knot.

shaded rectangular region and the top horizontal line of the rectangle is in the z-axis and the top hemi-spherical annuli are contained in different pages.

Using this definition, for a given link L, Furihata *et al.* [2] found an algorithm to find a flat plumbing basket surface from a closed braid $\overline{\beta} = L$. So we can define the *flat plumbing basket number* of L, denoted by $fpbk(L)$, to be the minimal number of flat annuli to obtain a flat plumbing basket surface of L.

Theorem 2.2. *([2]) Let L be an oriented link which is a closed n-braid with a braid word $\sigma_{n-1}\sigma_{n-2}\dots\sigma_1 W$ where the length of W is m and W has p positive letters, then there exists a flat plumbing basket surface S with $m+2p$ bands such that ∂S is isotopic to L, i.e., $fpbk(L) \leq m + 2p$.*

This upper bound has been improved by the third author [10] where the link is prime but not splittable.

Theorem 2.3. *([10]) Let L be an oriented link which is a closed n-braid with a braid word β whose length is m and let $ps(\sigma_i^{\pm 1})$ be the power sum of $\sigma_i^{\pm 1}$ in β for all $i = 1, 2, \dots, n-1$. Let γ be the cardinality of the set*

$$\Omega = \{i | 1 \leq i \leq n-1, \sigma_i \text{ and } \sigma_i^{-1} \text{ both appear in } \beta\}.$$

Let

$$\epsilon_i = \begin{cases} 1 & \text{if } 1 \leq ps(\sigma_i^1) \leq ps(\sigma_i^{-1}) \text{ or } ps(\sigma_i^{-1}) = 0, \\ -1 & \text{if } 1 \leq ps(\sigma_i^{-1}) \leq ps(\sigma_i^1) \text{ or } ps(\sigma_i^1) = 0. \end{cases}$$

Then the flat plumbing basket number of L is bounded by $m+n-1-4\gamma+2\sum_{i=1}^{n-1} ps(\sigma_i^{\epsilon_i})$, i.e.,

$$fpbk(L) \leq m + n - 1 - 4\gamma + 2\sum_{i=1}^{n-1} ps(\sigma_i^{\epsilon_i}).$$

The third author proved that every link L admits a flat plumbing basket representation from a canonical Seifert surface $\mathcal{F}_L$ of L to have a property that the Seifert graph $\Gamma(D_L)$ has a co-tree edge alternating spanning tree T [10].

Theorem 2.4. *([10]) Let Γ be an Seifert graph of canonical Seifert surface S of a link L with $|V(\Gamma)| = n$, $|E(\Gamma)| = m$ and the sign labeling ϕ. Let $G(\Gamma)$ be the Seifert*



FIGURE 3. (a) The knot 7_5 and (b) the numbering of 7_5 to find its Dowker-Thistlethwaite (DT)-code.

graph of Γ. *Let* T *be a co-tree edge alternating spanning tree of* Γ *and* μ *a labeling on* T *chosen in [10, Theorem 3.3]. Let* $\delta(T)$ *be the cardinality of the set*

$$\Psi(T) = \{e \in E(T) |\ \mu(e) \neq \phi(\overline{e}) \text{ for all } \overline{e} \in \Gamma(e)\},$$

and let $\zeta(T)$ *be the cardinality of the set*

$$\Upsilon(T) = \{\overline{e} \in E(\Gamma(T)) \mid \mu(e) = \phi(\overline{e}),\ \overline{e} \in \Gamma(e),\ e \in E(T) - \Psi(T)\}.$$

and let $\eta(T)$ *be the cardinality of the set*

$$\Phi(T) = \{\overline{e} \in E(\Gamma) - E(\Gamma(T)) \mid \mu(\overline{e}) = \nu(e)\}$$

where $\nu(e) = +(-,$ *resp.) if there is one extra positive(negative, respectively) sign in the path* P_e *joining end vertices of the edge* e *in* T. *Then the flat plumbing basket number of* L *is bounded by* $m - 3(n-1) + 2(2\delta(T) + \zeta(T) + \eta(T))$, *i.e.,*

$$fpbk(L) \leq m - 3(n-1) + 2(2\delta(T) + \zeta(T) + \eta(T)).$$

Next, we explain the *DT Code* (DT after C. Dowker and M. Thistlethwaite [1]) of a knot K because we will use it to identify prime knots whose flat plumbing basket number is 6. It can be obtained as follows:

(1) Start walking along K in a fixed direction as indicated in Fig. 3 (a) and count every crossing you pass through. If K has n crossings and given that every crossing is visited twice, the count ends at $2n$. Label each crossing with the values of the counter when it is visited, though when labeling by an even number, take it with a minus sign if you are walking "over" the crossing.
(2) Every crossing is now labeled with two integers whose absolute values run from 1 to $2n$. It is easy to see that each crossing is labeled with one odd integer and one even integer. The DT code of K is the list of even integers paired with the odd integers $1, 3, 5, \ldots$, taken in this order with the following modification: if the label is an even number and the strand followed crosses over at the crossing, then change the sign on the label to be a negative.

(3) Thus for example the pairing for the knot in Fig. 3 (b) is given

$$\begin{pmatrix} 1 & 3 & 5 & 7 & 9 & 11 & 13 \\ 12 & 8 & 10 & 14 & 4 & 2 & 6 \end{pmatrix}$$

At last, the DT code for this labelling is the sequence 12 8 10 14 4 2 6.

From the given flat plumbing basket surface in Fig. 2 (a), we obtain its DT code -32 -14 -44 -22 -40 2 28 10 -48 18 36 6 -24 -42 -12 30 -16 -46 -20 34 4 26 8 38 by choosing the starting point at the upper-left corner of the rectangle and the clockwise orientation.

3. The flat plumbing basket codes and results

For a given link L, Furihata *et al.* [2] found an algorithm to find a flat plumbing basket surface from a closed braid $\overline{\beta} = L$ as follows.

`Algorithm` [2]

- *Step* 1. For a give link L, we find its braid representation β, the closed braid $\overline{\beta} = L$.
- *Step* 2. Apply the method in [2] to obtain a flat plumbing basket surface $\mathcal{F}$ which is obtained from a disc by successively plumbing flat annuli.

This algorithm can be demonstrated in the following Example 3.1.

Example 3.1. *A flat plumbing basket code of the knot* 5_2 *is* (1, 2, 3, 4, 5, 6, 4, 5, 1, 2, 3, 6).

Proof. For the knot 5_2, we first present it as a closed braid $\overline{\sigma_2\sigma_1^{-1}(\sigma_2)^{-3}\sigma_1^{-1}}$ on three strings as illustrated in Fig. 4 (a). Although theorem in [2] stated differently, one can choose any two generators of the Artin's braid group B_3 as stated in Theorem 2.3. We choose the first $\sigma_2\sigma_1^{-1}$ to have a disc $\mathcal{D}$ which is the union of three discs, bounded by three Seifert circles, joined by two half twisted bands presented by $\sigma_2\sigma_1^{-1}$ as indicated by the dashed purple line in Fig. 4 (a). Since the rest word $(\sigma_2)^{-3}\sigma_1^{-1}$ has the length 4 and $(\sigma_2)^{-3}$ has the different sign to σ_2. we need three flat plumbings. However σ_1^{-1} has the same sign to σ_1^{-1}, we first change the sign of half twisted band by adding two flat annuli as shown in Fig. 4 (b). Now we pick as starting point as indicated as a red dot in Fig. 4 (b). Then, we read the flat bands along the disc $\mathcal{D}$ in the direction as given in Fig. 4 (b). By isotoping the original disc $\mathcal{D}$ to a standard disc as depicted as the rectangular gray region in Fig. 4 (c), we obtain a flat plumbing basket surface $\mathcal{F}$ in Fig. 4 (c). By rewriting labels in the set $\{7, 8, \ldots, 12\}$ by one in the set $\{1, 2, \ldots, 6\}$ depend on how $1, 2, \ldots, 6$ are connected to $7, 8, \ldots, 12$, and by the rule that for annulus presented by i is in front of the annulus presented by j whenever $i > j$, we obtain the first 12-tuple (1, 2, 3, 4, 5, 6, 4, 5, 1, 2, 3, 6) which is the flat plumbing basket code of the flat plumbing basket surface $\mathcal{F}$. □

Let us remark that since two groups of annuli presented by $\{1, 2, 3\}$ and $\{4, 5\}$ do not involve each other, one may obtain a different flat plumbing basket code of the knot 5_2, (1, 2, 6, 1, 2, 3, 4, 5, 6, 3, 4, 5) by changing (4; 5) into (1; 2) and (1; 2; 3) into (3; 4; 5).



FIGURE 4. (a) The knot 5_2 as a closed braid, (b) Seifert surface of 5_2 to apply algorithm, (c) a flat plumbing basket surface of 5_2.

One may see that the direction and the initial page of the open book decomposition was not fixed. By moving the initial page from 1 to 2, the effect on the flat plumbing basket code is an action by the permutation $\sigma = \begin{pmatrix} 1 & 2 & \dots & n-1 & n \\ 2 & 3 & \dots & n & 1 \end{pmatrix}$. By reversing the direction of pages in the open book decomposition, the effect on the flat plumbing basket code is an action by the permutation $\tau = \begin{pmatrix} 1 & 2 & \dots & n-1 & n \\ n & n-1 & \dots & 2 & 1 \end{pmatrix}$. By summarizing these observations, we find the following theorem.

Theorem 3.2. *For a positive integer n,*

(1) *The number of the* $2n$*-tuple flat plumbing basket codes presenting the same link are divisible by* n*.*
(2) *The number of the* $2n$*-tuple flat plumbing basket codes presenting the same link are divisible by* 2*.*

Now we will provide some classification theorems of knots and links by the flat plumbing basket numbers. Without using a computer program, the third author was able to prove the following classification theorem of links of the flat plumbing basket number $0, 1, 2, 3$ and 4 by using slightly different presentations of the flat plumbing basket surfaces, known as *permutation presentations*.

Theorem 3.3. *([12])*

(1) *A link* L *has the flat plumbing basket number* 0 *if and only if* L *is the trivial knot.*
(2) *A link* L *has the flat plumbing basket number* 1 *if and only if* L *is the trivial link of two components.*
(3) *A link* L *has the flat plumbing basket number* 2 *if and only if* L *is the trivial link of three components.*
(4) *A link* L *has the flat plumbing basket number* 3 *if and only if* L *is either the trivial link of four components or the Hopf link which is denoted by* $L2a1$*.*
(5) *A link* L *has the flat plumbing basket number* 4 *if and only if* K *is either the trefoil knot, the figure eight knot,* $L2a1 \sqcup O$*,* $L2a1\#L2a1$*,* $L6a5$*, or the trivial link of five components.*

where $\sqcup$ *presents the disjoint union.*

It is fairly easy to see that the number of components of the link whose flat plumbing basket number n is always congruent to $n+1$ modulo 2. Thus, the flat plumbing basket number of a prime knot has to be an even integer. From Theorem 3.3, one can easily obtain the following corollary which addresses the classification of all prime knots of the flat plumbing basket number 0, 2 and 4.

Corollary 3.4. (1) *A prime knot* K *has the flat plumbing basket number* 0 *if and only if* L *is the trivial knot.*
(2) *There does not exist a prime knot* K *whose flat plumbing basket number* 2*.*
(3) *A knot* K *has the flat plumbing basket number* 4 *if and only if* K *is either the trefoil knot or the figure eight knot.*

Example 3.5. (1) *The flat plumbing basket number of the link* 4_1^2 *is* 5*.*
(2) *The flat plumbing basket number of the knots* 5_2 *is* 6*.*

Proof. Flat plumbing basket surfaces of the link 4_1^2 with five annuli are depicted in Fig. 5. By Theorem 3.3, the link 4_1^2 can not have the flat plumbing basket number less than 5. Therefore, the flat plumbing basket number of the link 4_1^2 must be 5.

A flat plumbing basket surface of the link 5_2 with six annuli are depicted in Fig. 4 and its flat plumbing basket code is given (1, 2, 3, 4, 5, 6, 4, 5, 1, 2, 3, 6) in Example 3.1. By Corollary 3.4, the knot 5_2 can not have the flat plumbing basket number less than 6. □



FIGURE 5. (a) The link 4_1^2, (b) a flat plumbing basket surface of the link 4_1^2 with 5 flat plumbings.



FIGURE 6. A Type I move which decreases the flat plumbing basket number by 2.

Theorem 3.6. *The prime knot K has the flat plumbing basket number 6 if and only if it is either 5_1, 5_2, 6_1, 6_2, 6_3, 7_6, 7_7, 8_1, 8_3, 8_{12}, 8_{20}, 8_{21}, 9_{42}, 9_{44}, 9_{46}, 9_{48}, 10_{132}, 10_{136}, 10_{137}, 10_{140}, $11n_{38}$, $12n_{462}$, $13n_{973}$, $14n_{17954}$, $15n_{45460}$ or $16n_{246032}$.*

Proof. To consider all flat plumbing basket surface with 6 annuli, we first count all possible flat plumbing basket codes of $\{1,1,2,2,3,3,4,4,5,5,6,6\}$. Since these are presented as circular shapes, we can fix the first elements to be 1. Thus, there are $\dfrac{11!}{2^6} = 7,484,400$ many such flat plumbing basket codes.

The second author has written $C+$ program which determine whether a given flat plumbing basket code produces a knot or a link with more than 1 components, admits a Type I move as depicted in Fig. 6, finds a DT-code of the given flat plumbing basket code and performs series of reductions which realize the Reidemeister move type I and II. Using the program, we find that $6,415,200$ of them present links and $1,069,200$ present knots. Among $1,069,200$ flat plumbing basket codes presenting knots, there are $874,080$ codes which admit a Type I move, these knots must have the flat plumbing basket number less than 6. It leaves us $195,120$ flat plumbing basket codes. The computer program "knotfinder" of `Knotscape` reduces these DT-codes to the standard DT-codes and finds $105,162$ codes presents the unknot and $2,268$ codes present composite knots. All remaining $87,690$ reduced DT-codes of the flat plumbing basket codes are identified by the computer program "knotscape" of `Knotscape`. The results are listed as in Table 1. □

Name of knot	A flat plumbing basket codes	Number of f.p.b.codes
3_1	12341234 123124356456	$20,274$
4_1	12431243 123124365465	$32,442$
5_1	123456123456	12
5_2	123124563456	$4,176$
6_1	123124653465	$17,982$
6_2	123461253465	$1,368$
6_3	123456123654	$1,908$
7_6	123456123564	432
7_7	123456124365	$1,404$
8_1	124631254635	576
8_3	125314625463	288
8_{12}	123456124653	576
8_{20}	123612546534	$1,440$
8_{21}	123461256345	144
9_{42}	123614235465	720
9_{44}	123615246534	$1,152$
9_{46}	123614534625	$1,296$
9_{48}	123456125634	24
10_{132}	123612564534	144
10_{136}	125631243564	144
10_{137}	124615346253	288
10_{140}	123615624534	144
$11n_{38}$	126154635423	144
$12n_{462}$	124361546253	144
$13n_{973}$	126415364253	144
$14n_{17954}$	135264135264	36
$15n_{45460}$	124635124635	216
$16n_{246032}$	136254136254	72
composite		$2,268$
unknot	123124563564	$105,162$

TABLE 1. Knots of the flat plumbing basket surfaces with 6 bands out of $195,120$ cases.

Let us remark that the $C+$ program is available at the first author's homepage : `http://sec.pusan.ac.kr/?page_id=1303`. While using "knotfinder" of `Knotscape`, some of DT-codes of flat plumbing basket codes were not identified since the program aborts by the time limit set up by the program which may run infinitely because it repeatedly uses Reidemeister moves on DT-code which may increase the length of DT-code. However, we are lucky enough that at least one of DT-codes of the flat plumbing

basket codes which produce the same knot by the action of $\sigma = \begin{pmatrix} 1 & 2 & \dots & n-1 & n \\ 2 & 3 & \dots & n & 1 \end{pmatrix}$ as stated in Theorem 3.2 (1) makes an output DT-code. Further, because the length of these output DT-codes were less than or equal to 16 and all prime knots up to 16 crossings are tabulated, we can find the exact knot names. One can also observe that the number of the flat plumbing basket codes for a knot in Table 1 are all divisible by 12 except 3_1, 4_1 and 6_1 because some of flat plumbing basket codes are invariant by the action of σ or τ.

Since the trefoil knot and the figure eight knot have the flat plumbing basket number 4, there are exactly 26 prime knots whose flat plumbing basket numbers are exactly 6. The results in [8] found the flat plumbing basket number of prime knots up to 9 crossings except 24 knots. Using Theorem 3.6, we find the following corollary.

Corollary 3.7. (1) *The flat plumbing basket number of knots* 7_2, 7_4 *and* 9_{45} *is* 8.
(2) *The flat plumbing basket number of knots* 8_1, 9_{44} *is* 6.
(3) *The flat plumbing basket number of knots* 9_2, 9_5 *and* 9_{35} *are either* 8 *or* 10.

4. Conclusion

Authors already have all required DT-codes for the flat plumbing basket codes of the flat plumbing basket number 8 and 10. But we are not able to find a complete table like Table 1. There are two difficulties arise for these cases 1) the computer program "knotfinder" of `Knotscape` does not have a complete list of DT-codes for prime knots of more than 16 crossings, 2) the number of DT-codes we have to deal with increases exponentially. For example, a text.file contains all DT-codes for the flat plumbing basket codes of the flat plumbing basket number 10 is more than 2-gigabytes already.

To solve these problems, authors are trying to write a new computer program which identify whether two DT-notations are the same and revise "knotfinder" of `Knotscape`.

Acknowledgments

The TEX macro package PSTricks [21] was essential for typesetting the equations and figures. The third author was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science and Technology(2012R1A1A2006225).

References

[1] C. Dowker and M Thistlethwaite, Classification of knot projections, *Topology Appl.* **16** (1983) 19–31.
[2] R. Furihata, M. Hirasawa and T. Kobayashi, Seifert surfaces in open books, and a new coding algorithm for links, *Bull. London Math. Soc.* **40** (2008) 405–414.
[3] D. Gabai, The Murasugi sum is a natural geometric operation, in: Low-Dimensional Topology (San Francisco, CA, USA, 1981), Amer. Math. Soc., Providence, RI, 1983, pp. 131–143.
[4] D. Gabai, *The Murasugi sum is a natural geometric operation II*, in: Combinatorial Methods in Topology and Algebraic Geometry (Rochester, NY, USA, 1982), Amer. Math. Soc., Providence, RI, 1985, pp. –100.

[5] J. Gross, D. Robbins and T. Tucker, Genus distribution for bouquets of circles, *J. Combin. Theory B. Soc.* **47** (1989) 292–306.
[6] J. Gross and T. Tucker, Topological graph theory, Wiley-Interscience Series in discrete Mathematics and Optimization, Wiley & Sons, New York, 1987.
[7] J. Harer, How to construct all fibered knots and links, *Topology* **21** (1982) 263–280.
[8] S. Hirose and Y. Nakashima, Seifert surfaces in open books, and pass moves on links, *J. Knot Theory Ramifications* **22** 1450021 (2014) 1–12.
[9] C. Hayashi and M. Wada, Constructing links by plumbing flat annuli, *J. Knot Theory Ramifications* **2** (1993) 427–429.
[10] D. Kim, Basket, flat plumbing and flat plumbing basket surfaces derived from induced graphs, preprint, arXiv:1108.1455.
[11] D. Kim, The boundaries of dipole graphs and the complete bipartite graphs $K_{2,n}$, *Honam. math. J.* **36** (2014) 399–415, arXiv:1302.3829.
[12] D. Kim, A classification of links of the flat plumbing basket numbers 4 or less, *Korean J. of Math.* **22** (2014) 253–264.
[13] D. Kim, Y. S. Kwon and J. Lee, Banded surfaces, banded links, band indices and genera of links, *J. Knot Theory Ramifications* **22** 1350035 (2013), 1–18, arXiv:1105.0059.
[14] T. Nakamura, On canonical genus of fibered knot, *J. Knot Theory Ramifications* **11** (2002), 341–352.
[15] T. Nakamura, Notes on braidzel surfaces for links, *Proc. of AMS* **135** (2007), 559–567.
[16] L. Rudolph, Quasipositive annuli (Constructions of quasipositive knots and links IV.), *J. Knot Theory Ramifications* **1** (1992) 451–466..
[17] L. Rudolph, Hopf plumbing, arborescent Seifert surfaces, baskets, espaliers, and homogeneous braids, *Topology Appl.* **116** (2001) 255–277.
[18] H. Seifert, Uber das Geschlecht von Knoten, *Math. Ann.* **110** (1934), 571–592.
[19] J. Stallings, Constructions of fibred knots and links, in: Algebraic and Geometric Topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, CA, 1976), Part 2, Amer. Math. Soc., Providence, RI, 1978, pp. 55–60.
[20] M. Thistlethwaite, `Knotscape`, available at `http://www.math.utk.edu/~morwen/knotscape.html`.
[21] T. Van Zandt. PSTricks: PostScript macros for generic TEX. Available at `ftp://ftp.princeton.edu/pub/tvz/`.

School of Computer Science & Engineering, Pusan National University, Pusan, 609-735 Korea
E-mail address: `choi.yuno@gmail.com`

Gyeonggi Science High School, Suwon, 440-800, Korea
E-mail address: `jyg9628@nate.com`

Department of Mathematics, Kyonggi University, Suwon, 443-760 Korea
E-mail address: `dongseok@kgu.ac.kr`